MAGRUDER MEDICAL GROUP

Raymond D. Fuller, M.D.

IF YOU ARE ON THE FOLLOWING MEDICATIONS

**they MUST be stopped prior to your surgery/procedure date.
If they are not stopped, your surgery will be cancelled.**

- Aspirin (Anacin, Ascriptin, Bayer, Bufferin, Ecotrin, Excedrin)- **Hold 5 days**
- Brilinta (Ticagrelor)- **Hold 5 days**
- Coumadin (Warfarin)- **Hold 5 days**
- Diclofenac (Voltaren, Voltaren XR)- **Hold 5 days**
- Effient (Prasugrel)- **Hold 5 days**
- Eliquis (Apixaban) **Hold 48 hours**
- Lovenox (Enoxaparin)- **Hold 5 days**
- Meloxicam (Mobic)- **Hold 5 days**
- Oxaprozin (Daypro)- **Hold 5 days**
- Plavix (Clopidogrel)- **Hold 5 days**
- Pletal (Cilostazol) - **Hold 5 days**
- Pradaxa (Dabigatran Etexilate) **Hold 48 hours**
- Xarelto (Rivaroxaban) **Hold 48-72 hours**
- Naltexone- **Hold 72 hours**
- Jardiance- **Hold 72 hours**
- Metformin- **Hold 24 hours**
- Adipex- **Hold 72 hours**

***If you are taking any of the following GLP one inhibitors:**

- Dulaglutide (Trulicity)
- Exenatide (Bydureon, Byetta)
- Liraglutide (Saxenda, Victoza)
- Lixisenatide (Adlyxin)
- Semaglutide (Ozempic, Wegovy, Rybelsus, Mounjaro)
- Tirzepatide

Please hold *Please call prescribing Doctor in regards to bridging anti-diabetic therapy to avoid hyperglycemia if necessary

Daily dosing- Hold on day of surgery
Weekly dosing- Hold at least 1 week prior to surgery